FACTORS AFFECTING AVIAN USE OF RIPENING SUNFLOWER FIELDS

DIONN A. SCHAAF¹, GEORGE M. LINZ², AND WILLIAM J. BLEIER¹

¹Department of Biological Sciences, North Dakota State University, Fargo, ND 58105 ²USDA-APHIS, Wildlife Services, Bismarck, ND 58501

Introduction

In 1990, North Dakota's breeding blackbird population was estimated at more than 2.3 million pairs (Nelms et al., 1994). Sunflower has become an important part of these birds' diet from July-October, and damage due to blackbirds has been estimated at \$4-11 million (Linz et al., 1993). This problem has led to the use of various methods to control blackbird damage. Concern has been raised, however, on how some of these control methods may be affecting nontarget birds. Thus, we initiated a study on the use of the sunflower fields by non-blackbird species.

Methods

During the fall of 2000, twelve fields in Stutsman and Barnes counties in east-central North Dakota were used to assess the presence of non-target birds. Ten census points were laid out in each field. Censusing began within 15 minutes of sunrise. A 6-foot stepladder was used to aid in seeing over the sunflowers. A two-minute quiet period was used to allow disturbance to settle after reaching the census point, followed by an eight-minute count period. All birds were identified to species, sex, and age if possible. The location of each bird detected was recorded, including habitat type, direction, and distance from the census point. Distance was determined to the nearest 5m within 25m, while distances beyond 25m were recorded as 25-50m or over 50m. All birds flying over the plots were also recorded. Vegetation data were collected around each census point. Aerial photos of the fields and the surrounding area were taken, and a GIS program was used to quantify the habitat in and around the study fields.

Results

Each of the 12 fields was visited 7 times from late August to early October, for a total of 84 field censuses. A total of 82 non-blackbird species was observed, of which 49 were observed in sunflowers or foraging in flight over sunflowers. Of those 49 species, 30 were seed-eaters (Figure 1). While non-blackbirds were present throughout the study, they reached their highest numbers in late September, during migration (Figures 2 and 3).

Initial exploration of the data using the habitat figures produced by GIS also suggested a possible relationship between the surrounding habitat and the species seen in some of the fields. Pearson correlation coefficients were calculated for number of birds seen within 25m of a census point, against each habitat type by its proportion of the total area within 0.5 miles of the field (Figure 4). Some less common species were lumped by foraging group (insect- or seed-eating).

Further work will be done using the census data, vegetation data, and habitat figures to determine factors affecting avian presence in the sunflowers.

Acknowledgements

This research was funded jointly by the National Wildlife Research Center, a unit within the Wildlife Services program of the United States Department of Agriculture, Animal and Plant Health Inspection Service, and the Department of Biological Sciences at North Dakota State University.

Literature Cited

Linz, G. M., R. A. Dolbeer, J. J. Hanzel, and L. E. Huffman. 1993. Controlling blackbird damage to sunflower and grain crops in the northern Great Plains. U.S. Dept. of Agric., Animal and Plant Health Inspection Service, Agriculture Information Bull. No. 679.

Nelms, C. O., W. J. Bleier, D. L. Otis, and G. M. Linz. 1994. Population estimates of breeding blackbirds in North Dakota, 1967, 1981-1982, and 1990. Am. Midl. Nat. 132:256-263.




Figure 1. Foraging groups as a percentage of total non-blackbirds seen in ripening sunflower fields




Figure 2. Mean and standard error by census for birds seen in the sunflowers within 25m of census points





Figure 3. Total number of birds seen within 25m of a census point during seven censuses. Census 1: August 22-August 30; Census 2: August 29-September 6; Census 3: September 4-September 12; Census 4: September 12-September 19; Census 5: September 18-September 27; Census 6: September 27-October 5; Census 7: October 4-October 11.



- ♦ Grass/hay:Insectivores p=0.003
- ▲ Beans:Non-sparrow Seed-eaters p=0.03
- Farmyards:American robin p=0.02
- ♦ Grass:Savannah sparrow p=0.01
- △ Grass/hay:Song sparrow p=0.03

- Roads:Insectivores p=0.02
- ♦ Grain:Clay-colored sparrow p=0.02
- △ Wetland:Myrtle warbler p=0.05
- ☐ Grass/hay:Myrtle warbler p=0.04
- Roads:Savannah sparrow p=0.01

Figure 4. Significant Pearson correlation coefficients calculated for birds seen within 25m of a point against habitat around the field (n=12)

Proceedings of the 24th

SUNFLOWER RESEARCH WORKSHOP

January 17-18, 2002

Sponsored by:
National
Sunflower
Association

The 2001 Sunflower Research Workshop, sponsored by the National Sunflower Association, took place on January 17 and 18, 2002, at the Ramada Plaza Suites, Fargo, ND. The workshop was very well attended and received by public and private researchers from the United States and Canada, as well as other interested parties.

This volume contains nearly all the presentations given at the 2001 workshop. Some of the papers are summarized or abstract form.

The National Sunflower Association would like to extend its appreciation to those presenting papers/posters at this annual Sunflower Research Workshop and to those who participated by their

attendance and questions. Special thanks are extended to the NSA Research Forum Planning Committee, Dr. Gary J. Brewer, NDSU, Dr. Laurence D. Charlet, USDA-ARS and Pat Duhigg, Seeds 2000. Thanks also to Gerald Seiler, USDA-ARS-NCSL, Burton Johnson, NDSU, and Bob Benson, Mycogen Seeds for their expertise in moderating the workshop sessions.

Questions regarding these proceedings may be directed to the National Sunflower Association, 4023 State Street, Bismarck, ND 58503.

Note: The papers in these proceedings should not be reprinted in part or in total without the expressed consent of the author(s) involved.

TABLE OF CONTENTS

INSECTS Sunflower Crop Survey in North Dakota and	Selectivity of PPO-Inhibiting Herbicides in Sunflower59
South Dakota1	Kirk A. Howatt, Ron F. Roach, and
Arthur Lamey, Max Dietrich and Martin Draper	Sam F. Tutt
Insect Damage in North and South Dakota	PRODUCTION
Sunflower Fields in 2001: Results from the	Crop Sequence Calculator ² , V.2 – A Revised
National Sunflower Association Crop	Computer Program to Assist Producers63
Survey12	J. M. Krupinsky, D. L. Tanaka, J. S.
Larry D. Charlet	Fehmi, S. D. Merrill, M. A. Liebig,
•	J. R. Hendrickson, J. D. Hanson, R. L.
Lygus Bug and Kernel Brown Spot in Confection	Anderson, D. Archer, J. Knodel,
Sunflower: Determination of Economic Injury	P.A. Glogoza, L.D. Charlet, S. Wright,
Levels and Susceptible Growth Stages, and	And R. E. Ries (Retired)
Control Potential with Insecticides20	
Larry Charlet	T. J. Micromix – Effect of Micro-Nutrient
•	Combination on NuSun Sunflowers67
Prospects for Managing Sunflower Midge30 Gary J. Brewer	Tom D. Johnson and Jesse R. Barthel
J. 200	Sunflower Response to Limited Irrigation
Development of Chemical Attractants for	In Wyoming70
Sunflower Pest Insects	Craig M. Alford and Stephen D. Miller
	Available Soil Water, Sunflower Canopy
	Development and Productivity73
DIOTA CEC DISEASES	Robert M. Aiken
Effective Seed Treatments for the Control of	
Sclerotinia Root Infection in Sunflower44	The Jefferson Institute Approach to Crop
Khalid Rashid and John Swanson	Diversification81
Telling Adminstrate Dome Division	James Quinn
Head Rot Screening Nursery48	•
Bob Henson, Tom Gulya and	Soil Water Use and Soil Residue Coverage by
Brandon Miller	Sunflower Compared to Other
Diamon Miner	Crops88
WEEDS	Stephen D. Merrill, Donald L. Tanaka,
Clearfield* Sunflowers: Performance and	Joseph M. Krupinsky, Mark A. Liebig,
Progress54	John R. Hendrickson, Jonathan D. Hanson,
Mark L. Dahmer and Gary M. Fellows	and Ronald E. Ries

	stions for Including Sunflowers in	Improving Blackbird Population Control with	
Semia	rid Rotations97	Targeted Baiting Programs: Biological	
	Randy Anderson and Don Tanaka	Considerations	148
		Richard S. Sawin, George M. Linz, and	
Accen	tance of Bird Shield© by Growers:	William J. Bleier	
A 200	1 Season Field Report104	•	
A 200	Leonard R. Askham	Landscape Effects on Breeding Blackbird	
	Leonald IC Ashiam	Abundance and Sunflower Damage in the	
TT 1:		Southern Drift Plains of North Dakota	153
Herbi	cide Research in Sunflower105	Ryan L. Wimberly, George M. Linz,	13.
	Richard K. Zollinger and Jerry Ries		
		William J. Bleier, and H. Jeffrey Homan	
		Contact Discount Date of Contact I	
	DING & GENETICS	Spring Dispersal Patterns of Red-Winged	
	Resistance of Two Sulfonylurea-Resistant	Blackbirds Staging in East-Central	
	wer Sources to Selected Als	South Dakota	155
Herbi	zides117	H. Jeffrey Homan, George M. Linz,	
	A. Fabie and J. F. Miller	Richard M. Engeman, and Linda B. Penry	
Impro	ving Oil Quality in Sunflower Using		
	ld Relatives123	BREEDING AND GENETICS	
	Gerald J. Seiler	Nuclear Vigor Restoration Genes in Cultivated	
	V	Sunflower that Restore the Vigor Reducing	
		Cytoplasmic Effects of Perennial Helianthus	
QUAL	ITY	Species	150
-	ting NuSun Hybrid Oleic Acid	C.C. Jan and Juan A. Ruso	
	ntration Through Early Sampling126	CC Jan abu Juan 12 1030	
Сопсе	J. F. Miller, D. A. Rehder and B. A. Vick		
	j. P. Willer, D. A. Reitier and B. A. Vick		
	•	DISEASES	
	_	Variation in Head Rot Reactions Among	
OTHE		Sunflower Hybrids	162
	of Sunflower Seed Feeding on Conjugated	M. A. Draper and K. R. Ruden	
	c Acid Concentration in Milk Fat of		
Lactat	ing Dairy Cows131	D / LIDIT LIGHT	
	David B. Carlson and Chung S. Park	Bacterial Biological Control for	
	·	Sclerotinia Head Rot in Sunflower	
		Robert W. Duncan, W. G. D. Fernando and	
POSTI	ERS PRESENTED	Khalid Y. Rashid	
RIRD	PREDATION	Decline of Weed Densities in Sunflower as	
		Affected by Multiple Tactics in a Three-Crop	
	Use of Roadside Habitat and Implications	Rotation	166
FOF CA	ttail Management135	Ron Ries, Don Tanaka and Randy	200
	Bryan D. Safratowich, George M. Linz,	Anderson	
	William J. Bleier, and Carina J. Lee	11111113011	
_		Swafferman Designation and Bland	
	Affecting Avian Use of Ripening Sunflower	Sunflower Desiccation and Plant	454
Fields	139	Dry-Down	171
	Dionn A. Schaaf, George M. Linz, and	Burton L. Johnson, Timothy D.	
	William J. Bleier	Larson, and Robert A. Henson	
	·		
Baiting	Blackbirds During Spring Migration	Effect of Maturity and Achene Location on	
	h Dakota143	Fatty Acid Composition	176
	George M. Linz, Amy E. Barras, Richard A.	Tim D. Larson, Burton L. Johnson and	
	Sawin, William J. Bleier, H. Jeffrey Homan,	Brady A. Vick	
	David L. Bergman, and Linda B. Penry	•	
	waves in religious, and initial D. I tilly		
		Sunflower Date of Planting Study in Western	
		North Dakota 3-year Summary	182
		Roger O. Ashley, Eric D. Eriksmoen,	~~
		and M. Bridget Whitney	
		and it blinger willing	